

## IMPORTANT: PLEASE READ

### PART III: CONSUMER INFORMATION

#### Pr APO-BICALUTAMIDE Bicalutamide

This leaflet is part III of a three-part “Product Monograph” published when APO-BICALUTAMIDE was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about APO-BICALUTAMIDE. Contact your doctor or pharmacist if you have any questions about the drug.

#### ABOUT THIS MEDICATION

##### What the medication is used for:

APO-BICALUTAMIDE is used in the treatment of advanced prostate cancer in combination with other drugs (LHRH analogues) which reduce the levels of androgens in the body or surgery.

##### What it does:

Androgens are male sex hormones within the body which can cause tumor growth within the prostate. APO-BICALUTAMIDE belongs to a group of medicines called non-steroidal anti-androgens. This means that Apo- Bicalutamide interferes with some of the actions of androgens to prevent the tumor from growing.

##### What are the Stages of Prostate Cancer:

- Localized disease – the early stages of disease when prostate cancer is confined to the prostate gland.
- Locally advanced disease – the disease progresses and the cancer spreads to other tissues within the pelvis.
- Advanced or metastatic disease – the disease progresses to other parts of the body.

The PSA (Prostate Specific Antigen) test is a simple blood test for a protein produced by the prostate (PSA). This test has helped in the detection of prostate cancer resulting in an increase in the number of men whose prostate cancer is detected at an early stage.

##### What are the Treatment Options for Localized Prostate Cancer:

The optimal treatment for a given individual will depend on the specific circumstances of his case. For localized disease, patients are usually offered one of the following:

- Surgery to remove the prostate
- Targeted radiotherapy to kill the cancer cells in the prostate
- No treatment initially (watchful waiting) whereby the patient is monitored until there are signs of progression before treatment is started.

##### When it should not be used:

- If you have early phase (localized) prostate cancer requiring watchful waiting.
- If you are allergic to bicalutamide or any of the nonmedicinal ingredients in APO-BICALUTAMIDE.
- APO-BICALUTAMIDE must not be taken by women, including pregnant women or mothers who are breast feeding their babies.
- APO-BICALUTAMIDE must not be given to children.

##### What the medicinal ingredient is:

The active ingredient in APO-BICALUTAMIDE is Bicalutamide

##### What the important nonmedicinal ingredients are:

APO-BICALUTAMIDE 50 mg tablets contains the following nonmedicinal ingredients: anhydrous lactose, colloidal silicon dioxide, croscarmellose sodium and magnesium stearate.

##### What dosage forms it comes in:

APO-BICALUTAMIDE comes in tablets containing 50 milligrams (mg) of bicalutamide as the active ingredient.

#### WARNINGS AND PRECAUTIONS

- Apo-Bicalutamide should only be prescribed by a doctor experienced with the treatment of prostate cancer.
- Apo-Bicalutamide 150mg/day dose should not be used.
- Apo-Bicalutamide may rarely be associated with liver failure.

Before you use APO-BICALUTAMIDE talk to your doctor or pharmacist if:

- You have previously had a reaction to taking APO-BICALUTAMIDE or any of the ingredients in the product (see What the

Important Nonmedicinal Ingredients Are). As mentioned earlier, your tablets contain lactose which may cause a problem in a small number of patients who are sensitive to them.

- You are suffering from any disorder or disease, which affects your liver.

If you go into hospital let the medical staff know you are taking APO-BICALUTAMIDE.

Your tablets are unlikely to adversely affect your ability to drive a car or to operate machinery.

### INTERACTIONS WITH THIS MEDICATION

Please inform your doctor if you are taking or have recently taken any other medicines, even those not prescribed.

- In particular if you are taking oral anti-coagulants (to prevent blood clots)

Please note that these statements may also apply to products used some time ago.

### PROPER USE OF THIS MEDICATION

Follow your doctor's instructions about when and how to take your tablets. Ask your doctor or pharmacist if you are not sure.

- The usual adult dose is 50 mg daily.
- Swallow the tablet(s) whole with a drink of water.
- Try to take your dose at the same time each day.
- Do not stop taking your tablets even if you are feeling well, unless your doctor tells you.

During the first few months of use, you may be monitored by your physician for signs of changes in your liver function. In approximately 2.0% of patients, such changes may lead to withdrawal of therapy.

If you experience a rise in PSA while taking APO-BICALUTAMIDE, your physician may discontinue APO-BICALUTAMIDE for several weeks in order to monitor your condition off treatment.

### Overdose:

If you take more than your normal dose contact your doctor or nearest hospital.

### Missed Dose:

You should take APO-BICALUTAMIDE as prescribed. However, if you miss a dose do not take an extra dose. Just resume your usual schedule.

### SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like all medicines, APO-BICALUTAMIDE 50 mg can have side effects.

Side effects that are very common (more than 10 in every 100 patients are likely to have them):

- tender or enlarged breast tissue
- hot flushes

Side effects that are common (1 to 10 in every 100 patients are likely to have them):

- nausea
- diarrhoea
- itching
- feeling weak

Side effects that are rare (1 to 10 in every 10,000 patients are likely to have them):

- vomiting
- dry skin

Occasionally APO-BICALUTAMIDE may be associated with changes in your blood, which may require your doctor to do certain blood tests.

### SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM

Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
<b>Uncommon (1 to 10 in every 1000 patients are likely to have them)</b>			
Serious breathlessness, or sudden worsening of breathlessness, possibly with a cough or fever. Some patients taking APO-BICALUTAMIDE 50 mg get an inflammation of the lungs called interstitial lung disease.		√	

**SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM**

Symptom / effect	Talk with your doctor or pharmacist		Stop taking drug and call your doctor or pharmacist
	Only if severe	In all cases	
Severe itching of the skin (with raised lumps) or swelling of the face, lips, tongue and/or throat, which may cause difficulty in swallowing		√	
Yellow skin and eyes (jaundice). These may be symptoms of liver damage.		√	

Do not be alarmed by this list of possible events. You may not have any of them.

Tell your doctor or pharmacist if you think you have any of these or any other problems with your tablets.

***This is not a complete list of side effects. For any unexpected effects while taking APO-BICALUTAMIDE, contact your doctor or pharmacist.***

**HOW TO STORE IT**

- Keep your tablets in the container they came in.
- Do not take your tablets after the expiry date on the container. Dispose of them in an appropriate way.
- Keep your tablets in a safe place where children cannot see or reach them. Your tablets could harm them.
- Keep your tablets at room temperature (15 to 30 °C). Protect from light.

REPORTING SUSPECTED SIDE EFFECTS

To monitor drug safety, Health Canada collects information on serious and unexpected effects of drugs. If you suspect you have had a serious or unexpected reaction to this drug you may notify Health Canada by:

toll-free telephone: 866-234-2345  
 toll-free fax 866-678-6789  
 By email: [cadrmc@hc-sc.gc.ca](mailto:cadrmc@hc-sc.gc.ca)

By regular mail:  
 Canadian Adverse Drug Reaction Monitoring Program (CADRMP)  
 Health Canada  
 Address Locator: 0201C2  
 Ottawa, ON K1A 1B9

*NOTE: Before contacting Health Canada, you should contact your physician or pharmacist.*

**MORE INFORMATION**

For more information, please contact your doctor, pharmacist or other healthcare professional.

This leaflet plus the full product monograph, prepared for health professionals, can be obtained by contacting DISpedia, Apotex's Drug Information Service at:

1-800-667-4708

This leaflet can also be found at:  
<http://www.apotex.ca/products>.

This leaflet was prepared by Apotex Inc., Toronto, Ontario, M9L 1T9.

Last revised: June 5, 2007